

Fractal correlations in the CfA2-South redshift survey

M. Joyce^{1,2,3} M. Montuori¹ and F. Sylos Labini^{1,4}

Received _____; accepted _____

Submitted to Astrophys. J. Letters

¹INFM Sezione Roma 1, Università di Roma "La Sapienza" P.le A. Moro 2, 00185 Roma, Italy

²INFN Sezione Roma 1, Università di Roma "La Sapienza" P.le A. Moro 2, 00185 Roma, Italy

³School of Mathematics, Trinity College, Dublin 2, Ireland.

⁴ Dépt. de Physique Théorique, Université de Genève, 24, Quai E. Ansermet, CH-1211 Genève, Switzerland

ABSTRACT

We report our analysis of the properties of galaxy clustering for a new redshift sample of galaxies, the CfA2-South catalog, using statistical methods which do not rely on the assumption of homogeneity. We find that, up to $\sim 20h^{-1}Mpc$, which is the largest scale to which correlation properties can be reliably inferred, the galaxy clustering is scale-invariant and characterized by a fractal dimension $D = 1.9 \pm 0.1$. Further there is no statistical evidence for homogeneity at any of the larger scales (up to $\sim 150h^{-1}Mpc$) probed more weakly by the catalog. These results means that characteristic “correlation lengths” for the clustering of galaxies derived using standards methods of analysis are not meaningful. Further the results are very consistent with those obtained from many other catalogs using the methods adopted here, which show the $D \approx 2$ fractal continuing to beyond $100h^{-1}Mpc$. The incompleteness of the relevant data conjectured by various authors to give rise to such behaviour is therefore proved to have no significant effect (up to $20h^{-1}Mpc$) on the measured correlations.

Subject headings: galaxies: general; galaxies: statistics; cosmology: large-scale structure of the universe

1. Introduction

That the galaxy distribution exhibits fractal properties has been well established for almost twenty years (Mandelbrot, 1977, Peebles, 1980). Controversy still surrounds questions concerning the range of the fractal regime, the value of the fractal dimension and the eventual presence of a cross-over to homogeneity (Pietronero et al., 1997, Davis 1997, Wu et al., 1998 and references therein). In this letter we present the answers to these questions provided by a statistical analysis of a new large redshift sample of galaxies, the CfA2-South survey (Huchra et al., 1998).

Red-shift catalogs such like this one have been, and continue to be, extensively analyzed with the standard two-point correlation function $\xi(r)$ method (Peebles, 1980) which involves normalizing to a mean density extracted from the catalog. Indeed the predictions of all standard type theories of structure formation are usually framed in terms of these statistics (or related ones such as the power spectrum). One result of this analysis is that up to a small scale ($\sim 10 \div 20h^{-1}Mpc$) power-law type behaviour is observed in $\xi(r)$ which is interpreted to indicate fractal behaviour with a characteristic fractal dimension $D \approx 1.3$. The central result however is the derivation of a characteristic length, the “correlation length”, $r_0 \approx 5h^{-1}Mpc$, which should mark the end of a fractal behaviour and the onset of a trend towards homogenization: It is defined as the scale at which the the density fluctuations become of the same order as the average density, and at a few times this distance the fluctuations have small amplitude i.e the density tends to some well-defined asymptotic homogeneous value. This trend to homogeneity at such a scale is in apparent agreement with the structure-less angular data, but it is puzzling with respect to the much larger structures observed in the 3-d data. From this perspective it seems that the presence or absence of structures in the data is irrelevant for the determination of r_0 .

Puzzled by this apparent anomaly Pietronero and collaborators (Pietronero

1987, Coleman & Pietronero 1992, Sylos Labini et al., 1998) reconsidered the analysis of galaxy correlations with such methods, adopting instead a procedure in which the correlations are characterized with statistics which are appropriate whether there is homogeneity or not, and which allow the underlying assumption of homogeneity in the standard analysis to be tested. The conclusion of such considerations applied to various galaxy surveys such as CfA1, SSRS1, IRAS, APM-Stromlo, Perseus-Pisces and LEDA (Sylos Labini et al., 1998) is that there is in fact no statistical evidence for the assumed homogeneity on the scales probed by any of these catalogs (up to $\sim 100h^{-1}Mpc$ for the LEDA catalog), and that the derived “correlation length” therefore has no meaning except in relation to the particular sample. On the other hand there is indeed clear fractal behaviour at the scales probed by all these catalogs, but it is correctly characterized by a fractal dimension $D \approx 2$ rather than the smaller value derived in the standard analysis. These results have been questioned on the basis of the statistical validity of the catalogs (Davis 1997). Various authors have in fact proposed that the incompleteness of these data may lead to an apparent fractal behaviour. In this context the catalog we analyse in this paper is important as it is a new and very accurate one, which does not suffer from the possible problems indicated with other surveys. Indeed it has been analysed extensively using the standard methods (Park et al., 1994) and is frequently used as a constraint on theories of structure formation.

Rather than being tested directly with methods such as those used here, the assumption of homogeneity is often defended with the assertion that the “correlation length” is indeed a real physical scale characterizing the clustering of galaxies, because it shows the stability in different samples indicative of homogeneity and does not show the behaviour which would be associated with a fractal distribution. In practice, however, correlation lengths are not observed to be very stable. For example in the SSRS2 catalog Benoist et al. (1996) have measured different values of r_0 in the range $[4, 15]h^{-1}Mpc$. In the CfA catalog as

analysed by Park et al., 1994 there is also a large variation of the measured correlation length, which we will discuss below. It is here that concept of “luminosity selection bias” enters (Davis et al., 1988, Park et al., 1994, Benoist et al., 1996): Galaxies of different brightness are supposed to be clustered differently, and it is this which is proposed as the physical explanation of the observed variation, rather than the absence of real underlying homogeneity. Rather than there being one real scale characterizing the correlations of galaxies, there is then an undetermined number of such scales. In our discussion below we will pay particular attention to this point, showing that the variation of the “correlation length” in the analysis of Park et al. is perfectly consistent with our results, and that the account of luminosity selection bias invoked to explain it is therefore unnecessary.

We first describe our methods of analysis before proceeding to the catalog and our results. Essentially we make use of a very simple statistic (Pietronero 1987, Coleman & Pietronero 1992, Sylos Labini et al., 1998) which is appropriate for characterizing the properties of regular as well as irregular distributions. This is the conditional average density, defined as

$$\Gamma(r) = \left\langle \frac{1}{S(r)} \frac{dN(< r)}{dr} \right\rangle \quad (1)$$

where $dN(< r)$ is the number of points in a shell of radius dr at distance r from an occupied point and $S(r)dr$ is the volume of the shell. The average indicated by the angle brackets is over all the occupied points contained in the sample. Another related quantity we use is

$$\Gamma^*(r) = \frac{3}{4\pi r^3} \int_0^r \Gamma(\rho) d^3\rho, \quad (2)$$

which is just the conditional density in a sphere of radius r rather than in a spherical shell as for $\Gamma(r)$ ⁵. It is thus a more stable global quantity which smoothes out the rapidly varying fluctuations which may appear in $\Gamma(r)$.

⁵Note that in real samples the integral in eq.2 has a lower cut-off different from zero.

These statistics are suitable both for the study of the approach (if any) to homogeneity of a given distribution, as well as for the identification of fractal properties. The former is indicated by the tendency to a constant as a function of distance r , while the latter will be indicated by a simple power-law behaviour. Specifically for a scale invariant distribution of points with fractal dimension D we have:

$$\Gamma(r) = \frac{BD}{4\pi} r^{-\gamma} \quad \text{and} \quad \Gamma^*(r) = \frac{3B}{4\pi} r^{-\gamma} \quad (3)$$

where $\gamma = 3 - D$ and B is the lower cut-off, characterizing the end of self-similarity in the finite set. (B is the number of galaxies contained in a ball of radius $1h^{-1}Mpc$ around a given galaxy.)

In the analysis of the correlation properties of a real system with these statistics there are two important physical scales defining the range in which one can infer real correlation information from them:

(i) The upper cut-off R_s up to which the statistic can be calculated. It is simply the size of the largest sphere around any galaxy which can be inscribed inside the sample volume, since the conditional density is computed in complete shells (Coleman & Pietronero 1992, Sylos Labini et al., 1998). It depends on the survey geometry - on the solid angle of the survey and the effective depth of the particular sub-sample we analyse. As one approaches this scale both the number of points being averaged over and the separation of these points drops, so that one is no longer computing a real average correlation. We will return to this point below in discussing our results.

(ii) A lower cut-off $\langle\Lambda\rangle$, which is related to the number of points contained in the sample. It is simply the scale below which the behavior of the conditional density is dominated by the sparseness of the points. In $\Gamma(r)$ this regime is characterized by highly fluctuating behaviour, while in the $\Gamma^*(r)$ it is manifest as $1/r^3$ decay away from any finite value.

The problems of the standard analysis can readily be seen from the fact that, for the case of a fractal distribution, the standard “correlation function” $\xi(r)$ and “correlation length ” r_o are just

$$\xi(r) = \frac{\Gamma(r)}{\langle n \rangle} - 1 \quad \text{and} \quad r_o = \left(\frac{BD}{8\pi \langle n \rangle} \right)^{\frac{1}{\gamma}} \quad (4)$$

where $\langle n \rangle$ is the average density in the sample as estimated in the calculation of $\xi(r)$. In a fractal $\langle n \rangle$, and therefore r_o , are quantities which characterize not intrinsic properties of the distribution, but the particular sample one is considering. For example, in a sample of depth R_s the density is, on average, $\langle n \rangle = \Gamma^*(R_s)$. Thus for a $D = 2$ fractal, the *average* value in such samples is $r_o = R_s/3$. Further, for the case of a fractal, $\xi(r)$ is a power law only for $r \ll r_o$. The deviation from this behaviour at large scales is again just due to the size of the observational sample and does not correspond to any real change of the correlation properties. The log derivative of eq.4 with respect to $\log(r)$ is

$$\gamma' = \frac{d(\log(\xi(r)))}{d \log(r)} = -\frac{2\gamma r_o^\gamma r^{-\gamma}}{2r_o^\gamma r^{-\gamma} - 1} \quad (5)$$

where r_o is defined by $\xi(r_o) = 1$. The tangent to $\xi(r)$ at $r = r_o$ has a slope $\gamma' = -2\gamma$. It is thus clear that, even if the distribution has fractal properties, it is very difficult to recover the correct slope from the study of the $\xi(r)$ function.

We now turn to the data. The CfA2 South galaxy sample (Huchra et al., 1998) contains 4390 galaxies with magnitude $m_{B(0)} \leq 15.5$ covering $20^h \leq \alpha \leq 4^h$ in right ascension and $-2.5^\circ \leq \delta \leq 90^\circ$ in declination. This part of the sky includes regions where the galactic extinction has been measured to be important, which needs to be corrected for. For the purposes of comparison of our results with the analysis carried out by Park et al., 1994 we perform these corrections in exactly the same way as these authors, by excluding the same regions: $20^h \leq \alpha \leq 21^h$; $3^h \leq \alpha \leq 4^h$; $21^h \leq \alpha \leq 2^h$ and $b > -25^\circ$ and $2^h \leq \alpha \leq 3^h$ and $b > -45^\circ$ where b is the galactic latitude. The catalog contains a large supercluster called

the *Perseus-Pisces chain* and it reveals large voids in the foreground and background of this supercluster. The complex galaxy structures found in this catalog have been described by Huchra et al., 1998. Note that all the wide angle redshift surveys (i.e. CfA1, SSRS1, SSRS2, Perseus-Pisces) have shown comparable fluctuations, which indeed represent the subject of our analysis.

A redshift survey limited in this way by apparent magnitude has a systematic selection effect: At large distances only the brighter galaxies are included in the sample, while at small distances there are also the fainter ones. To avoid this selection bias it is a standard procedure to consider the so-called volume-limited (VL) samples (e.g. Davis & Peebles, 1983), in which one chooses an upper cut-off in distance and takes only all the galaxies which are bright enough to be seen up to this distance. In Tab.1 we report the characteristics of the VL samples we have considered, characterized by successive limits in absolute magnitude with a step of 0.5 magn. The distances have been computed from the linear Hubble law ($r = cz/H_0$ and $H_0 = 100h \text{ km/sec/Mpc}$ where $0.5 < h < 1$) and the absolute magnitude from the standard relation $M = m - 5 \log_{10} r(1 + z) - 25 - Kz$, where for the K correction we have taken $K = 3$ (as in Park et al., 1994). The results of our analysis are extremely weakly dependent on this latter correction, as they are on modifications to the $r - z$ relation corresponding to different cosmological models, simply because the maximum red-shift is so small ($z < 0.05$).

In fig.1 the conditional average density $\Gamma^*(r)$ is plotted for each of these five samples. The error bars displayed correspond to the variance on 20 bootstrap resamplings of each sample⁶. The behaviour of these errors can also be understood in terms of the two limits we discussed above: At small distances they are large because the results for the average

⁶For a sample with N points, a random integer is generated between 1 and N for each point. Those points with the same number are discarded to produce the sub-sample.

quantities are dominated by a few points which make the average density non-zero. In some cases they are large at $r \sim R_s$ because here one is averaging about just a few well separated galaxies, one of which is removed in the resampling. It is important to note however that although the behaviour of these errors is related to the two cut-offs, they *cannot* be taken to be measurement errors for these effects, which are systematic. At small scales the real behaviour of the conditional density is determined by the real lower cut-off and is intrinsically highly fluctuating no matter how well sampled; at large scales no resampling of the points around a single galaxy can tell us what the intrinsic variance is in the quantity measured from different independent points.

An examination of the figure shows that beyond a scale, which grows with the depth of the VL sample, there is in each case a well defined power-law, until a scale near to the upper cut-off R_s at which, in some samples, it shows a deviation towards a flatter behaviour. The first scale is just the lower cut-off $\langle \Lambda \rangle$ due to sparseness discussed above. It can be checked quantitatively that its increase in the deeper samples scales with the growing mean distance between points. To perform a fit to these curves we also need to take account of the systematic effect as one approaches $r \sim R_s$, not included in the bootstrap errors, due to the non-averaging. In principle we cannot know how large the error at this scale is since we do not know the real variance in the density at this scale. The criterion we use here to place an upper cut-off up to which we assume this systematic effect is not important is a simple one: We require that the average distance between the centres of the spheres we are averaging over at the relevant depth r be such that the spheres do not overlap. The quantitative meaning of this can be read off from the figure inserted in figure one, as the point where the average distance becomes equal to twice the depth. Obviously this scale R_u grows with sample size and we see it reaches a maximum of about $20h^{-1}Mpc$ in our deepest VL sample.

EDITOR: PLACE FIGURE 1 HERE.

In each of the sample we perform a best fit to the dimension D in the range of scales $[\langle\Lambda\rangle, R_u]$ (see Tab.1). Our result is that the dimension is $D = 1.9 \pm 0.1$ in this range of scales probed by the samples i.e. from $0.5h^{-1}Mpc$ to $20h^{-1}Mpc$.

The normalization of the conditional average density in different VL samples depends on the luminosity selection function of the sample considered. The procedure to perform such a normalization has been described in Sylos Labini et al., 1998. In short we assume that the joint luminosity and space density can be written as

$$\nu(M, r) = \Gamma(r) \cdot \phi(M) = \frac{DB}{4\pi} r^{D-3} \cdot \phi(M) \quad (6)$$

where the luminosity function $\phi(M)$ has been normalized to unity ($\int_{-\infty}^{M_{min}} \phi(M) dM = 1$ and M_{lim} is the faintest absolute magnitude contained in the sample). We can associate to a VL sample limited at M_{VL} a luminosity factor:

$$\Phi(M_{VL}) = \int_{-\infty}^{M_{VL}} \phi(M) dM . \quad (7)$$

The normalization of the space density is then

$$\Gamma_{norm}(r) = \frac{\Gamma(r)}{\Phi(M_{VL})} = \frac{DB}{4\pi} r^{D-3} \cdot \frac{1}{\Phi(M_{VL})} \quad (8)$$

From eq.8, we can estimate the parameter B of the distribution. We find $B \approx (12 \pm 2h^{-1}Mpc)^{-D}$, which agrees very well with the value found in various other catalogs (Sylos Labini et al., 1998). This concordance clearly shows that the results derived in these catalogs (which are in good agreement with each other) are not significantly effected by incompleteness or other imperfections, at least at the scales probed by the present analysis. The dismissal on these grounds of such very robust and self-consistent results thus seems (even at larger scales) rather inappropriate.

Beyond $\sim 20h^{-1}Mpc$ we cannot reliably infer average properties of the galaxy distribution from the CfA2 South catalog alone. We can see however that, at least to $\sim 30h^{-1}Mpc$, there is evidence of a clearly fluctuating behaviour consistent with the continuation of the fractal. We can further examine unaveraged quantities like the number counts $N(< r)$ from the origin. These show the highly fluctuating behaviour typical of a fractal, with slopes ranging from 3.5 to 2 (corresponding to a global overdensity on the scale of the survey). Are we potentially missing evidence for a real flattening in the conditional density by assuming that the variance is large i.e could the deviations at $r \sim R_s$ from the simple power-law in a few samples be real averaged ones? Could the slopes closer to 3 in a few samples in the number counts be real average behaviour? One simple way in which one can discount this possibility is to show that the particular behaviour is not an average one, even for the sample, by showing it to be associated with a particular feature within the sample. In the present case the evidence for this is very clear-cut: These behaviours are all associated with the main overdensity in the sample, the Perseus-Pisces supercluster. The judgement one might make at first glance, that this catalog is shows no tendency to homogeneity simply because of this feature, is quite simply correct.

Finally we return to the standard $\xi(r)$ analysis, in particular as applied by Park et al., 1994 to the CfA2 catalog. Our findings of power-law correlations without evidence for a cut-off imply that the normalization to a mean density to derive a “correlation length” is *conceptually* flawed. Calculationally, however, there is nothing wrong with deriving such a scale, and the results should be perfectly consistent (numerically) with those given here. In Table 1 we list the simple volume density $\langle n \rangle$ in each of the VL samples. Using this as our normalizing density in equation (4), and the values of B and D from the measured $\Gamma^*(r)$, we obtain r_o in the range $5 \div 10h^{-1}Mpc$. These values are very compatible with those which can be read off from Figure 10 in Park et al.(for the whole CfA survey) which gives values for r_o in the VL samples shown of approximately $7h^{-1}Mpc$ and $9h^{-1}Mpc$. As

we discussed above the *average* r_o predicted in a continuing $D = 2$ fractal for such samples is $r_o = R_{VL}/3$. The ratio of the measured r_o therefore agrees strikingly well with this average scaling since the two samples have respective depths $101h^{-1}Mpc$ and $130h^{-1}Mpc$. The difference between the absolute values for r_o and the *average* r_o implies that the CfA2 survey would correspond in this case to an overdensity by a factor of four (relative to the average density $\Gamma^*(R_{VL})$ at the depth of the survey). To invoke (as Park et al., 1994 do) the additional hypothesis of luminosity selection bias to explain all the variation of the observed r_o is in our view very problematic unless one has first clarified the role of the intrinsic variance in the densities to which one is normalizing to obtain r_o .

We conclude by noting again that our results on this survey permit only the establishment of a fractal with $D \approx 2$ to $\sim 20h^{-1}Mpc$. While we find no statistical evidence in this survey for the hypothesis of homogeneity up to the largest scales probed ($\sim 150h^{-1}Mpc$), we cannot exclude that the fluctuations we see in non-averaged quantities rule out any trend to homogeneity at such scales. The question of the true properties of the galaxy distribution at such scales will only be definitively clarified with the advent of the much larger forthcoming red-shift surveys (2dF and SLOAN). The application of methods such as those used here, which do not assume homogeneity, is essential for the resolution of this fundamental observational question.

We warmly thank L. Pietronero for useful discussions and collaborations. This work has been partially supported by the EEC TMR Network “Fractal structures and self-organization” ERBFMRXCT980183 and by the Swiss NSF.

REFERENCES

- Benoist C., *et al.* 1996 *Astrophys. J.* **472**, 452-459
- Coleman, P.H. and Pietronero, L., 1992 *Phys.Rep.* **231**, 311-391
- Davis, M., 1997 p.50-60 in the Proc. of the Conference “*Critical Dialogues in Cosmology*”
N. Turok Ed. World Scientific
- Davis, M., Peebles, P. J. E. 1983 *Astrophys. J.*, **267**, 465-482
- Davis M. *et al.*, 1988 *Astrophys. J. Lett.* **333**, L9-L12
- Huchra J.P., Vogeley M.S. and Geller M.J. 1998 To Appear in *Astrophys.J. Suppl.*
- Kaiser N., 1984 *Astrophys.J.* **284**, L9-L12
- Mandelbrot, B.B., 1977 “*Fractals: Form, Chance and Dimension*”, W.H.Freedman
- Park C., Vogeley M., Geller M. and Huchra J., 1994 *Astrophys. J.* **431**, 569-585
- Peebles, P.E.J., 1980 “*The Large Scale Structure of The Universe*” Princeton Univ.Press.
- Peebles P. J. E., 1993 “*Principles of physical cosmology*”, Princeton Univ. Press
- Pietronero L., 1987 *Physica A*, **144**, 257-284
- Pietronero, L., Montuori, M. and Sylos Labini, F., 1997 p.24-49 in the Proc of the
Conference “*Critical Dialogues in Cosmology*” N. Turok Ed. World Scientific
- Sylos Labini F., Montuori M., Pietronero L., 1998 *Phys.Rep.* **293**, 66-226
- Sylos Labini F. and Montuori M., 1998 *Astronom.Astrophys.* **331**, 809-814
- Wu K.K.S., Lahav O. and Rees M., 1998 *Nature* submitted (astro-ph/9804062)

Sample	R_{VL}	M_{VL}	N	R_s	$\langle n \rangle$	$\langle \Lambda \rangle$	R_u	D
VL185	60.2	-18.5	724	13.0	$1.3 \cdot 10^{-2}$	0.5	7	2.0
VL19	74.9	-19.0	622	18.0	$6.0 \cdot 10^{-3}$	0.7	8	1.8
VL195	92.9	-19.5	520	18.5	$2.5 \cdot 10^{-3}$	1.2	12	1.9
VL20	115	-20.0	292	24.0	$7.5 \cdot 10^{-4}$	3.0	15	1.9
VL205	141.7	-20.5	132	29.6	$2.0 \cdot 10^{-4}$	6.0	20	1.8

Table 1: Volume limited samples of the CfA2-South. R_{VL} ($h^{-1}Mpc$) is the depth of the VL sample, M_{VL} is the absolute magnitude limit of the VL sample, N the number of galaxies contained, $\langle n \rangle = N/V$ where $V = (\Omega/3)R_{VL}^3$, R_s ($h^{-1}Mpc$) is the effective depth and D is the estimated fractal dimension in the range $[\langle \Lambda \rangle, R_u]$.

Fig. 1.— The conditional average density $\Gamma^*(r)$ in the different VL samples of the CfA2-South galaxy sample. The error bars displayed correspond to the variance measured for 20 bootstrap resampling of each sample. The values of the power law fit are reported in Tab.1. In the inserted panel we show the average distance between the centers of the spheres (see text): the solid line corresponds to the curve $\langle d \rangle = 2r$ where $\langle d \rangle$ is the average separation between the the centers of the spheres and r is the depth.

